

Durham Research Online

Deposited in DRO:

11 June 2019

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Felde, Vivian A. and Flantua, Suzette G. A. and Jenks, Cathy R. and Benito, Blas M. and de Beaulieu, Jacques-Louis and Kuneš, Petr and Magri, Donatella and Nalepka, Dorota and Risebrobakken, Bjørge and ter Braak, Cajo J. F. and Allen, Judy R. M. and Granoszewski, Wojciech and Helms, Karin F. and Huntley, Brian and Kondratien, Ona and Kalnia, Laimdota and Kupryjanowicz, Mirosława and Malkiewicz, Małgorzata and Milner, Alice M. and Nita, Małgorzata and Noryskiewicz, Bożena and Pidek, Irena A. and Reille, Maurice and Salonen, J. Sakari and Šeirien, Vaida and Winter, Hanna and Tzedakis, Polychronis C. and Birks, H. John B. (2020) 'Compositional turnover and variation in Eemian pollen sequences in Europe.', *Vegetation history and archaeobotany*, 29 (1). pp. 101-109.

Further information on publisher's website:

<https://doi.org/10.1007/s00334-019-00726-5>

Publisher's copyright statement:

This is a post-peer-review, pre-copyedit version of an article published in *Vegetation History and Archaeobotany*. The final authenticated version is available online at: <https://doi.org/10.1007/s00334-019-00726-5>

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Compositional turnover and variation in Eemian pollen sequences in Europe

Vivian A. Felde^{1,23,*}, Suzette G. A. Flantua¹, Cathy R. Jenks¹, Blas M. Benito¹, Jacques-Louis de Beaulieu², Petr Kuneš³, Donatella Magri⁴, Dorota Nalepka⁵, Bjørg Risebrobakken⁶, Cajo J. F. ter Braak⁷, Judy R. M. Allen⁸, Wojciech Granoszewski⁹, Karin F. Helmens¹⁰, Brian Huntley⁸, Ona Kondratienė¹¹, Laimdota Kalniņa¹², Mirosława Kupryjanowicz¹³, Małgorzata Malkiewicz¹⁴, Alice M. Milner¹⁵, Małgorzata Nita¹⁶, Bożena Noryśkiewicz¹⁷, Irena A. Pidek¹⁸, Maurice Reille², J. Sakari Salonen¹⁹, Vaida Šeirienė²⁰, Hanna Winter²¹, Polychronis C. Tzedakis^{22,#}, H. John B. Birks^{1,22,23,#}

¹ Department of Biological Sciences, University of Bergen, PO Box 7803, N-5020 Bergen, Norway

² Aix Marseille Université, Avignon Université, CNRS, IRD, IMBE, Aix-en-Provence, France

³ Department of Botany, Faculty of Science, Charles University, Prague, Czech Republic

⁴ Department of Environmental Biology, Sapienza University of Rome, Italy

⁵ Department of Palaeobotany, W. Szafer Institute of Botany, Polish Academy of Sciences, Lubicz 46, 31-512 Kraków, Poland

⁶ NORCE Climate and Bjerknes Centre for Climate Research, Jahnebakken 5, N-5007 Bergen, Norway

⁷ Biometris, Wageningen University and Research, Wageningen, The Netherlands

⁸ Department of Biosciences, Durham University, Durham, DH1 3LE, UK

⁹ Polish Geological Institute – National Research Institute, Carpathian Branch, Skrzatów 1, 31-560 Kraków, Poland

¹⁰ Department of Physical Geography and the Bolin Centre for Climate Research, Stockholm University, 106 91 Stockholm, Sweden

¹¹ Nature Research Centre, Institute of Geology and Geography, T. Ševčenkos 13, 03223 Vilnius, Lithuania

¹² Faculty of Geography and Earth Sciences, University of Latvia, Rainis Boulevard 19, LV-1586 Riga, Latvia

¹³ Department of Palaeobotany, Institute of Biology, University of Białystok, Ciołkowskiego 1J, 15-425 Białystok, Poland

¹⁴ Laboratory of Paleobotany, Department of Stratigraphic Geology, Institute of Geological Sciences, University of Wrocław, Cybulskiego 34, 50-205 Wrocław, Poland

¹⁵ Department of Geography, Royal Holloway, University of London, Egham, TW20 0EX, UK

16 Department of Fundamental Geology, Faculty of Earth Sciences, University of Silesia, Będzińska 60, 41-200
Sosnowiec, Poland

17 Faculty of Earth Sciences, Nicolaus Copernicus University in Toruń, Lwowska 1, 87-100 Toruń, Poland

18 Faculty of Earth Sciences and Spatial Management, M. Curie Skłodowska University, al. Krasnicka 2cd, 20-
718 Lublin, Poland

19 Department of Geosciences and Geography, University of Helsinki, PO Box 64, FI-00014 Helsinki, Finland

20 Nature Research Centre, Institute of Geology and Geography, T. Ševčenkos 13, 03223 Vilnius, Lithuania

21 Polish Geological Institute and National Research Institute, Rakowiecka 4, 00-975 Warszawa, Poland

22 Environmental Change Research Centre, University College London, London, WC1E 6BT, UK

23 Bjerknes Centre for Climate Research, University of Bergen, Norway

*** Author for correspondence:** Vivian A. Felde (vivian.felde@uib.no)

H. John B. Birks and Polychronis C. Tzedakis contributed equally to this paper and are considered
joint senior authors

[Pending – to include ORCID]

Vivian A. Felde [ORCID 0000-0002-2855-0894], Suzette G. A. Flantua, Cathy R. Jenks, Blas M.
Benito, Jacques-Louis de Beaulieu, Petr Kuneš, Donatella Magri, Dorota Nalepka, Bjørge
Risebrobakken, Cajo J. F. ter Braak, Judy R. M. Allen, Wojciech Granoszewski, Karin F. Helms,
Brian Huntley, Ona Kondratienė, Laimdota Kalniņa, Mirosława Kupryjanowicz, Małgorzata
Malkiewicz, Alice M. Milner, Małgorzata Nita, Bożena Noryśkiewicz, Irena A. Pidek, Maurice Reille, J.
Sakari Salonen, Vaida Šeirienė, Hanna Winter, Polychronis C. Tzedakis, H. John B. Birks

Abstract

The Eemian interglacial represents a natural experiment on how past vegetation with negligible human impact responded to amplified temperature changes compared to the Holocene. Here, we assemble 47 carefully selected Eemian pollen sequences from Europe to explore geographical patterns of i) total compositional turnover and total variation for each sequence and ii) stratigraphical turnover between samples within each sequence using detrended canonical correspondence analysis, multivariate regression trees, and principal curves. Our synthesis shows that turnover and variation are highest in central Europe (47–55°N), low in southern Europe (south of 45°N), and lowest in the north (above 60°N). These results provide a basis for developing hypotheses about causes of vegetation change during the Eemian and their possible drivers.

62

63 **Keywords:** detrended canonical correspondence analysis, extrinsic processes, inertia, intrinsic
64 processes, last interglacial dataset, multivariate regression trees, neutral processes, principal curves

Introduction

The last interglacial (Eemian, c. 129–116 thousand years ago (ka)) is the most extensively studied pre-Holocene stage of the Quaternary (Tzedakis 2007a). It is characterised during its early part by a stronger summer insolation anomaly, peak global mean surface air temperatures of $\sim 1^{\circ}\text{C}$ above pre-industrial values, reaching 3–11 $^{\circ}\text{C}$ in the Arctic (Fischer et al. 2018), and by a peak sea-level of 6–9 m above present (Dutton et al. 2015).

The duration of the last interglacial represents the interval of reduced ice volume, demarcated at its onset by deglaciation and at its close by glacial inception (Tzedakis et al. 2012) and is broadly equivalent to Marine Isotope sub-Stage 5e (~ 132 –116 ka) and the Eemian interglacial of north-west Europe (Kukla et al. 2002). The term Eemian was introduced by Harting (1874) to describe a subsoil characterised by warm marine molluscs in the Eem valley near Amersfoort in The Netherlands (Tzedakis 2007a). Integrated palaeoceanographic and pollen analyses from the Portuguese Margin show that the marine isotopic and terrestrial stage boundaries are not synchronous, with the interval of temperate forest conditions extending from ~ 129 to ~ 111 ka (Shackleton et al. 2003; Tzedakis et al. 2018). Here we use the term ‘Eemian’ informally to refer to the forested interval (protocratic, mesocratic, and oligocratic/telocratic phases – see Fig. 1a) in last interglacial pollen sequences across Europe. While the long duration ($\sim 18,000$ years) of the forest interval in southern Europe is supported by independent chronologies (Brauer et al. 2007), a shorter duration ($\sim 11,000$ years) has generally been applied to north-central European pollen sequences on the basis of a partially annually laminated record at Bispingen, Germany at 53 $^{\circ}\text{N}$ (Müller 1974). However, recent joint palaeoceanographic pollen analyses from the Bay of Biscay (Sánchez Goñi et al. 2012) and comparisons with pollen sequences in southern France and southern Germany indicate a long Eemian duration ($\sim 18,000$ years) at least as far north as 48 $^{\circ}\text{N}$. It is possible that the unlaminated upper part of the Bispingen sequence represents a longer time interval than presently assumed and that the duration of the forested interval in northern Germany was approximately as long as farther south. In the absence of any independent chronology and duration estimates, the length of the Eemian in Fennoscandia remains unclear.

Ever since the pioneering studies in Denmark and Germany by Jessen and Milthers (1928), many Eemian pollen sequences have been analysed, focussing mainly on sedimentary settings, stratigraphies, pollen assemblages, vegetation histories, dating, and climate (e.g. Zagwijn 1996; Kühl 2003; Tzedakis 2007a). Eemian pollen records present a valuable opportunity to study vegetation patterns across broad spatial and temporal scales without extensive human impact (Kühl 2003; Tzedakis 2007a; Milner et al. 2013). Ecological questions abound about the Eemian. For example, are Eemian inferred vegetation patterns similar to Holocene patterns prior to human influence? How similar are Eemian pollen stratigraphies across Europe? Are there consistent temporal patterns in the appearance, expansion, and decline of major arboreal taxa across Europe? What are the spatial variations in pollen compositional change (“turnover”) and total variation across Europe?

To answer such questions and to study vegetation patterns and trends during the Eemian in Europe, we compile an Eemian dataset based on 47 representative pollen sequences. Here, we consider what the spatial patterns are in pollen turnover and total variation across Europe. We use compositional turnover and variation to estimate change in pollen-assemblage composition over time and space (Andersen 1994; Birks and Birks 2004; Birks 2007). With these explorative analyses presented here, we address the following two questions. **Q1)** How does compositional turnover change within Eemian pollen sequences? **Q2)** What are the spatial variations in total pollen compositional turnover and total variation across Europe?

As the concept of turnover is rarely used in pollen analysis (see Birks 2007), we summarise what this concept is. In community ecology, turnover is used to describe and possibly to quantify the replacement of one species by another in an assemblage in space or time or both. In pollen analysis, turnover is used to refer to the amount of compositional change of all pollen taxa within a stratigraphical sequence, namely along a temporal gradient (Birks 2007). Although the concept of turnover is widely used in ecology (e.g. Baselga 2010; Descombes et al. 2017) and biogeography (e.g. Buckley and Jetz 2008), little appears to be known about how turnover within an assemblage changes with time (e.g. Jarzyna et al. 2014). Pollen sequences provide a means of studying biotic turnover over long time periods. We use compositional turnover (one type of β -diversity *sensu* Anderson et al. (2011)) as estimates of change in pollen-assemblage composition along the temporal gradient in a pollen sequence. This is “directional turnover” (Anderson et al. 2011) or “compositional gradient length” (Tuomisto 2010). We avoid referring to β -diversity because it now has so many meanings (e.g. Tuomisto 2010; Anderson et al. 2011) in ecology, biogeography, and palaeoecology.

Based on previous studies of vegetation development during interglacials spanning part of one precession cycle (e.g. Andersen 1994; Birks and Birks 2004; Tzedakis 2007b; Helmens 2014), we expect all sequences to show a generally unimodal pattern of compositional change or turnover within the Eemian reflecting the protocratic, mesocratic, and oligocratic plus telocratic phases (**Q1**; Fig. 1a). Additionally, we predict the total amount of turnover to have been highest in northern Europe, intermediate in central, and lowest in southern Europe, whereas total variation may have been highest in central Europe and lowest in the north (**Q2**; Fig. 1b). These hypotheses are based on the assumption that at a different mix of extrinsic, intrinsic, and neutral processes play out in the different regions. At the beginning of the Eemian in the south, most taxa would have already been present and compositional change was likely driven primarily by extrinsic and/or intrinsic processes (*sensu* Williams et al. 2011a), such as climatic shifts, competition, and facilitation, along with some neutral processes such as historical legacies and location of glacial-stage refugia (Jackson and Blois 2015). In the north, turnover may reflect species spread as driven by extrinsic and neutral processes. In the mesocratic phase, changes in all regions may have been a result primarily of intrinsic and neutral processes (e.g. Iversen 1960; Birks 1986) with some extrinsic processes, whereas changes in the

oligocratic/telocratic phase may have been driven by an interaction of extrinsic, intrinsic, and neutral processes.

Dataset and methods

Answering the two questions above (Q1, Q2) requires three major components – i) representative Eemian pollen data across Europe, ii) critical screening to ensure they lack pre- or post-Eemian pollen spectra or hiatuses, are of comparable analytical standard, and have a consistent pollen nomenclature, and iii) robust numerical tools for consistent data analysis. Here we describe the methodology followed to address these components. See Electronic Supplementary Materials (ESM) 1 for details of the numerical methods and software.

Compiling a European Eemian pollen dataset

We implemented four criteria (Box 1) to select Eemian pollen sequences from different sources, including Pangaea, the European Pollen Database, and the Polish Pleistocene Pollen Database (Kupryjanowicz et al. 2018a) (see ESM 2 for details). For sequences in geographically critical areas with few complete Eemian sequences and the primary data no longer available, published Eemian diagrams were digitised. Pollen values are expressed as percentages of total pollen excluding pollen of aquatics and all spores.

Box 1.

Criteria for including sequences in the Eemian dataset

1. Must cover the entire Eemian and display signals of protocratic, mesocratic, and oligocratic/telocratic phases of an interglacial, allowing for differences in how these phases are reflected in different parts of Europe (Birks 1986)
2. Have at least 15 analysed samples
3. Have no clear evidence of major hiatuses
4. Have consistent pollen identifications of reasonable analytical standard

Box 2.

Assumptions (see also ESM 3)

Palynological

- i) Sequences are all Eemian, are complete with no discernible hiatuses, and have roughly constant monotonic sediment accumulation rates
- ii) A minimum number of 15 samples in a sequence is adequate to assess variability and turnover. Range of sample numbers is 16–213 (mean = 60; median = 52)
- iii) Pollen taxonomy is of a comparable and consistent standard for all sequences, with all major non-arboreal taxa identified and recorded for each sequence. Range of taxa is 13–99 (mean = 48; median = 47)
- iv) Digitised data are complete and of sufficient quality for analysis
- v) In the absence of independent chronologies from northern Europe, we assume that the duration of the forested interval across Europe was approximately similar

Numerical

- vi) DCCA provides robust estimates of turnover and variation (see also ESM 1)
- vii) Multivariate regression trees (MRT) and associated cross-validation are robust in identifying the optimal

partition tree, even with small sample numbers (Simpson and Birks 2012) and hence in accessing the amount of palynological variation in a sequence

viii) Numerical results are not overly affected by pollen-count size and hence number of taxa

ix) Numerical results are not overly affected by the number of samples in a sequence

Because samples in sequences are in stratigraphical order, numerical analysis should, when appropriate, take account of this data-property and be constrained by sample order. Ideally, this constraint should be sample age but age estimates are not available for almost all Eemian sequences. In the absence of age estimates, we have used depth which reflects sample order within a sequence. Birks (2007) discusses using either age or depth as external constraints in the ordination of Holocene sequences and obtains almost identical results irrespective of the type of constraint imposed. The palynological and numerical assumptions behind our study are summarised in Box 2 and are discussed more fully in ESM 3.

Estimating compositional turnover within each sequence and total turnover for each pollen sequence

Compositional turnover is estimated using detrended canonical correspondence analysis (DCCA; constrained by depth (or order) plus depth²) (Birks 2007; ter Braak and Šmilauer 2012) and principal curves (PCs) (Simpson and Birks 2012).

DCCA directly scales variables' (in our case pollen taxa) ordination scores such that their average within-sample standard deviation is unity along the ordination axes which are here a mix of sample depth or order. The change in weighted average (WA) sample scores (*CaseR sensu* ter Braak and Šmilauer 2012) reflects compositional change or turnover in standard deviation (SD) units. PCs are more "neutral" than DCCA in that they make fewer assumptions of the data than DCCA does. In the PC approach, a PC is fitted to the entire Eemian dataset of 2840 samples. Sample locations along the final PC are determined and scaled to 0–1. Maximum difference of sample scores within a sequence is a relative turnover measure (Simpson and Birks 2012). For each sequence, total compositional turnover is estimated and within each sequence we explore patterns of turnover. Emphasis here is placed on the DCCA results for both total turnover and changes within a sequence because they are expressed in ecologically interpretable units of standard deviation (SD) of taxon turnover (Figs. 2–3; ESM 8). The PC results for total turnover are summarised in Fig. 3b.

Estimating total variation for a sequence

Two contrasting numerical approaches are used to estimate total variation for each sequence: 1) total inertia (=weighted variance) estimated by correspondence analysis for each sequence in the absence of any external constraints (ter Braak and Šmilauer 2012) and 2) sequence partitioning using multivariate regression trees (MRTs) (Simpson and Birks 2012) and cross-validation to estimate the optimal number of partitions ("zone boundaries"). Optimal partition number depends on the number of samples in a sequence. We thus express the number of optimal partitions as a proportion of the

number of samples. This reflects the amount of palynological variation within a sequence. Estimates of variation are summarised in Fig. 3c and d.

Results

We assemble 47 suitable sequences from 14 countries grouped into three regions (above 60°N, 45–60°N, below 45°N; Fig. 2; ESM 2). The sequences range from 16–213 samples and 13–99 pollen taxa after basic taxonomic harmonisation (ESM 4). The dataset is chosen to cover as much of Europe as possible and to contain representative (but not all) Eemian sequences from Europe. Some areas, however, have many more Eemian sequences than others (e.g. N Germany and Poland compared with Scandinavia).

There is a distinct temporal pattern of compositional change within each sequence over geographical space displayed by the individual sample scores (SD units) on DCCA axis 1 (Figs. 2–3; ESM 8). The observed patterns indicate site-specific changes, but it is also possible to identify two general patterns where the DCCA sample values show either a gradually changing linear trend or a unimodal pattern where it reaches a peak and then slowly declines. Of the 31 sequences with a unimodal pattern, 27 occur in central Europe. Four northern sequences show a unimodal pattern and no southern sequence shows such a pattern. Linear trends are found in all regions (3 in northern, 8 in central, 5 in southern Europe) (Fig. 2; ESM 8).

The greatest Eemian turnover and variation are in central Europe, whereas the southern and northern regions show less change (Fig. 3; ESM 9). Total compositional turnover (DCCA, Fig. 3a; PC, Fig 3b) and inertia (Fig. 3c) have peaks between 47 and 55°N. Lowest values are above 60°N and are low south of 45°N. In contrast, the proportion of optimal partitions has highest values in central and north Europe (Fig. 3d). The null hypothesis that the mean change in optimal partitions does not differ between regions is not rejected. An analysis of variance indicates a probability >0.05 that the null hypothesis is true ($F = 2.575$, $p = 0.088$). Full analytical results are given in ESM 4–7.

Discussion

We compiled an Eemian pollen sequence dataset from across Europe, based on a set of predefined criteria and assumptions, to explore the magnitude of compositional turnover and total variation in Eemian pollen sequences. We show that there is substantial spatial variation in both variables, presumably reflecting responsiveness to various degrees of extrinsic, intrinsic, and neutral processes at local and regional scales during the Eemian (Birks 1986; Williams et al. 2011a; Jackson and Blois 2015).

Our Eemian dataset shows that there are important geographical voids of complete Eemian pollen sequences – none in Ireland and Iberia, few in Britain and France, and, not surprisingly, very few in much of Scandinavia that was heavily glaciated after the Eemian. Studies focusing on interglacial

dynamics in comparison to the Holocene would benefit from a better geographical coverage of Eemian sequences to assess all vegetation types and climate conditions of the continent.

Pollen sequences in central Europe display the expected unimodal pattern (Fig. 2), but surprisingly, several sequences from across Europe, especially in the north and the south, show a gradually changing linear trend (cf. Fig. 1b). Less change in the south and the north (Fig. 3) could imply that the taxa are shifting abundances between a few equally abundant taxa, thereby displaying a gradual changing pattern of turnover instead of a unimodal pattern (Q1).

We show that the greatest Eemian variation and turnover are in central Europe (Fig. 3), whereas there are fewer compositional changes in the southern and the northern regions (Q2). This is only partly consistent with the prior expectation of a unimodal trend in turnover during the entire Eemian (Fig. 1b). One possible explanation for the greatest change in central Europe is that during the Eemian there was a great variety and replacement of forest trees and shrubs (e.g. *Carpinus*, *Corylus*, *Quercus*, *Taxus*, *Tilia*, *Ulmus*) contributing to the regional pollen deposition, thereby providing a greater potential for palynological compositional turnover to be detected by a technique such as DCCA. Such pollen taxa have high N2 values where N2 (ter Braak and Verdonschot 1995) is the effective number of occurrences of a given taxon. DCCA is based on weighted averages, and N2 is effectively determined by taxa with high abundances (ter Braak and Verdonschot 1995).

An alternative and more convincing hypothesis is that in southern Europe many taxa were already present at the onset of the Eemian (Bennett et al. 1991; Tzedakis et al. 2013), and the observed pollen-stratigraphical changes there may be mostly a result of intrinsic processes such as facilitation and competition between species and of neutral processes leading to a gradual shift from a landscape dominated by evergreen sclerophyll and deciduous-*Quercus* and *Ulmus* to a vegetation with later expansion of *Carpinus*, *Ostrya*, and *Abies* into locally favourable habitats. Such changes may be recorded palynologically as relatively low turnover (Fig. 3).

An additional related hypothesis for the observed contrasting patterns in southern and central Europe (Fig. 3) involves a detailed consideration of the pollen records. While a number of southern European pollen sequences show a pattern of early, middle, and late expanding taxa not dissimilar to that of central Europe (Tzedakis et al. 2001), the main taxa (e.g. deciduous-*Quercus*, *Ulmus*) tend to persist through most of the Eemian with later expansion of *Carpinus*, *Ostrya*, and *Abies* superimposed on the existing assemblages. In central Europe, by comparison, there is a replacement of the early dominant taxa by later arrivals (similar to a relay), leading to a more accentuated and greater turnover. These patterns (relay vs expansion plus persistence of early taxa) may reflect climate differences. As climate shifted towards cooler temperatures during the course of the Eemian (Fig. 1a), the early thermophilous taxa may have declined in central Europe, while in the south, temperature (and precipitation) may not have become limiting, allowing the persistence of these taxa (e.g. Bennett et al. 1991), resulting in reduced turnover and variation.

The low palynological turnover and variation in northern Europe (Fig. 3) may reflect the relatively low pollen richness with few abundant taxa of restricted competitive abilities, especially trees, in the Eemian. The restricted arboreal flora may simply have been a result of climatic limitations. The detailed study at Sokli in northern Finland (Salonen et al. 2018) suggests, however, that thermophilous taxa such as *Corylus* were present and persisted in the north until the close of the Eemian, perhaps because of a major decrease in seasonality in the late Eemian with increase in winter insolation.

An alternative general hypothesis for the observed patterns in turnover is “silent palynological turnover”. North of the Alps today there are only two native *Quercus* species, one native *Pinus* species, one native *Abies* species, and one native *Juniperus* species. In contrast, in southern Europe all of these genera comprise several, if not many (e.g. *Quercus*), species. The pollen, however, of the various species within these genera cannot generally be distinguished. This pollen species-morphological limitation may result in compositional turnover at the species level not being detected. In central and northern Europe with only one or two species in these genera, turnover is at or near the species level and is thus more visible palynologically. “Silent turnover” could thus contribute, at least in part, to the observed patterns within Europe as a whole (Fig. 3). It could also suggest the apparent persistence of some genera in southern European sequences and result in a potentially misleading record of the amount of ecological turnover that actually took place during the Eemian.

These contrasting hypotheses as explanations for the patterns detected in our study (Fig. 3) highlight how much there is to be discovered about the palaeoecology and vegetation dynamics of the Eemian.

The DCCA compositional turnover for the Eemian is higher than for the Holocene (11 ka) in over 40 sites in Scandinavia and Britain (Birks, unpublished data; Birks 2007). Future studies include extending the geographical coverage of Holocene turnover estimates into central and southern Europe so as to compare Eemian and Holocene turnover for nearby sites.

Conclusions

We have compiled a dataset of 47 representative Eemian pollen sequences from across Europe. We have consistently analysed the sequences using multivariate numerical methods to estimate total pollen compositional change (turnover) and total variation for each sequence and to explore how composition turnover changes within each sequence and between sequences. The turnover and variation estimates show coherent but unexpected geographical and temporal patterns. They provide a basis for developing hypotheses about palynological changes during the Eemian and their possible causes. This demonstration that numerical analysis of our Eemian dataset produces robust and ecologically interpretable patterns gives confidence in our dataset. Further questions about Eemian vegetation dynamics and history will be explored using this dataset in subsequent studies.

Data sources. See ESM 2.

Author contributions. Ideas – VAF, HJBB, BB; data compilation, digitising – SGAF, CRJ; data management, harmonisation, screening – VAF, BB, HJBB, SGAF; computing – VAF, BB; discussion and/or detailed comments –J-LdB, WG, BH, PK, DM, BR, CJFtB; writing and editing – HJBB, VAF, PCT, SGAF, CRJ; data contributions –JRMA, J-LdB, WG, KH, BH, LK, OK, PK, MK, MM, AMM, MN, BN, IAP, MR, JSS, VS, PCT, HW

All authors read and commented on the manuscript, and gave final approval for publication.

Acknowledgements and funding We thank Prof. Konrad Wolowski for granting us access to the Polish Pleistocene Pollen Database. We are also very grateful to the European Pollen Database (www.europeanpollendatabase.net/) and the work of the EPD data contributors and the EPD community for making EPD data publicly available. HJBB is indebted to Hilary Birks for many valuable discussions. HJBB, SGAF, and CRJ are supported by the ERC Advanced Grant 741413 Humans on Planet Earth (HOPE). VAF is supported by IGNEX-eco (6166) funded by VISTA - a basic research program in collaboration between The Norwegian Academy of Science and Letters, and Equinor; BB is supported by NFR project IGNEX (249894). This paper is a contribution to the IGNEX and IGNEX-eco projects.

References

- Andersen ST (1994) History of the terrestrial environment in the Quaternary of Denmark. *Bulletin of the Geological Society of Denmark* 41:219-228
- Anderson MJ, Crist TO, Chase JM, Vellend M, Inouye BD, Freestone AL, Sanders NJ, Cornell HV, Comita LS, Davies KF, Harrison SP, Kraft NJB, Stegen JC, Swenson NG (2011) Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist. *Ecology Letters* 14:19-28. doi:10.1111/j.1461-0248.2010.01552.x
- Baselga A (2010) Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography* 19 (1):134-143. doi:10.1111/j.1466-8238.2009.00490.x
- Bennett KD, Tzedakis PC, Willis KJ (1991) Quaternary refugia of north European trees. *Journal of Biogeography* 18 (1):103-115
- Birks HJB (1986) Late Quaternary biotic changes in terrestrial and limnic environments, with particular reference to north west Europe. In: Berglund BE (ed) *Handbook of Holocene Palaeoecology and Palaeohydrology* J Wiley & Sons, Chichester, pp 3-65
- Birks HJB (2007) Estimating the amount of compositional change in late-Quaternary pollen-stratigraphical data. *Vegetation History and Archaeobotany* 16:197-202. doi:10.1007/s00334-006-0079-1
- Birks HJB, Birks HH (2004) The rise and fall of forests. *Science* 305 (5683):484-485
- Brauer A, Allen JRM, Mingram J, Dulski P, Wulf S, Huntley B (2007) Evidence for last interglacial chronology and environmental change from southern Europe. *Proceedings of the National Academy of Sciences USA* 104:450-455
- Buckley LB, Jetz W (2008) Linking global turnover of species and environments. *Proceedings of the National Academy of Sciences USA* 105 (46):17836-17841. doi:10.1073/pnas.0803524105

- Descombes P, Vittoz P, Guisan A, Pellissier L (2017) Uneven rate of plant turnover along elevation in grasslands. *Alpine Botany* 127 (1):53-63. doi:10.1007/s00035-016-0173-7
- Dutton A, Carlson AE, Long AJ, Milne GA, Clark PU, DeConto R, Horton BP, Rahmstorf S, Raymo ME (2015) Sea-level rise due to polar ice-sheet mass loss during past warm periods. *Science* 349 (6244):aaa4019. doi:10.1126/science.aaa4019
- Fischer H, Meissner KJ, Mix AC, Abram NJ, Austermann J, Brovkin V, Capron E, Colombaroli D, Danianu A-L, Dyez KA, Felis T, Finkelstein SA, Jaccard SL, McClymont EL, Rovere A, Sutter J, Wolff EW, Affolter S, Bakker P, Ballesteros-Cánovas JA, Barbante C, Caley T, Carlson AE, Churakova O, Cortese G, Cumming BF, Davis BAS, de Vernal A, Emile-Geay J, Fritz SC, Gierz P, Gottschalk J, Holloway MD, Joos F, Kucera M, Loutre M-F, Lunt DJ, Marcisz K, Marlon JR, Martinez P, Masson-Delmotte V, Nehrbass-Ahles C, Otto-Bliesner BL, Raible CC, Risebrobakken B, Sánchez Goñi MF, Arrigo JS, Sarnthein M, Sjolte J, Stocker TF, Velasquez Álvarez PA, Tinner W, Valdes PJ, Vogel H, Wanner H, Yan Q, Yu Z, Ziegler M, Zhou L (2018) Palaeoclimate constraints on the impact of 2 °C anthropogenic warming and beyond. *Nature Geoscience* 11 (7):474-485. doi:10.1038/s41561-018-0146-0
- Harting P (1874) De bodem van het Eemdal. Verslagen en Mededelingen van de Koninklijke Academie van Wetenschappen, Afdeling Naturkunde II 8:282-290
- Helmens KF (2014) The Last Interglacial-Glacial cycle (MIS 5-2) re-examined based on long proxy records from central and northern Europe. *Quaternary Science Reviews* 86:115-143. doi:10.1016/j.quascirev.2013.12.012
- Iversen J (1960) Problems of the early post-glacial forest development in Denmark. *Danmarks Geologiske Undersøgelse IV* 4 (3):1-32
- Jackson ST, Blois JL (2015) Community ecology in a changing environment: Perspectives from the Quaternary. *Proceedings of the National Academy of Sciences USA* 112 (16):4915-1921. doi:10.1073/pnas.1403664111
- Jarzyna MA, Finley AO, Porter WF, Maurer BA, Beier CM, Zuckerberg B (2014) Accounting for the space-varying nature of the relationships between temporal community turnover and the environment. *Ecography* 37 (11):1073-1083. doi:10.1111/ecog.00747
- Jessen K, Milthers V (1928) Stratigraphical and paleontological studies of interglacial fresh-water deposits in Jutland and northwest Germany. *Danmarks Geologiske Undersøgelse II* 48:1-379
- Kühl N (2003) Der Bestimmung botanisch-klimatologischer Transferfunktion und die Rekonstruktion der hodennahen Klimazustandes in Europa während der Eem-Warmzeit. *Dissertationes Botanicae* 375:1-149
- Kukla GJ, Bender ML, de Beaulieu J-L, Bond G, Broecker WS, Cleveringa P, Gavin JE, Herbert TD, Imbrie J, Jouzel J, Keigwin LD, Knudsen K-L, McManus JF, Merkt J, Muhs DR, Müller H, Poore RZ, Porter SC, Seret G, Shackleton NJ, Turner C, Tzedakis PC, Winograd IJ (2002) Last Interglacial Climates. *Quaternary Research* 58 (1):2-13. doi:10.1006/qres.2001.2316
- Kupryjanowicz M, Nalepka D, Pidek IA, Walanus A, Balwierz Z, Bińka K, Fiłoc M, Granoszewski W, Kołaczek P, Majecka A, Malkiewicz M, Nita M, Noryśkiewicz B, Winter H (2018a) The east-west migration of trees during the Eemian Interglacial registered on isopollen maps of Poland. *Quaternary International* 467:178-191. doi:10.1016/j.quaint.2017.08.034
- Milner AM, Müller UC, Roucoux KH, Collier REL, Pross J, Kalaitzidis S, Christanis K, Tzedakis PC (2013) Environmental variability during the Last Interglacial: a new high-resolution pollen record from Tenaghi Philippon, Greece. *Journal of Quaternary Science* 28 (2):113-117. doi:10.1002/jqs.2617
- Müller H (1974) Pollenanalytische Untersuchungen und Jahresschichtenzählungen an der holsteinzeitlichen Kieselgur von Munster-Breloh. *Geologisches Jahrbuch A21*:107-140
- Salonen JS, Helmens KF, Brendryen J, Kuosmanen N, Välranta M, Goring S, Korpela M, Kylander M, Philip A, Pliik A, Renssen H, Luoto M (2018) Abrupt high-latitude climate events and decoupled seasonal trends during the Eemian. *Nature Communications* 9 (1):#2851. doi:10.1038/s41467-018-05314-1

- Sánchez Goñi MF, Bakker P, Desprat S, Carlson AE, van Meerbeeck CJ, Peyron O, Naughton F, Fletcher WJ, Eynaud F, Rossignol L, Renssen H (2012) European climate optimum and enhanced Greenland melt during the Last Interglacial. *Geology* 40 (7):627-630. doi:10.1130/G32908.1
- Shackleton NJ, Sánchez Goñi MF, Pailler D, Lancelot Y (2003) Marine Isotope Substage 5e and the Eemian interglacial. *Global and Planetary Change* 36 (3):151-155. doi:10.1016/S0921-8181(02)00181-9
- Simpson GL, Birks HJB (2012) Statistical learning in palaeolimnology. In: Birks HJB, Lotter AF, Juggins S, Smol JP (eds) *Tracking Environmental Change Using Lake Sediments Volume 5: Data Handling and Numerical Techniques*. Springer, Dordrecht, pp 249-327. doi:10.1007/978-94-007-2745-8_9
- ter Braak CJF, Šmilauer P (2012) *Canoco Reference Manual and User's Guide: software for ordination (version 5.0)*. Microcomputer Power, Ithaca, New York
- ter Braak CJF, Verdonschot PFM (1995) Canonical correspondence analysis and related multivariate methods in aquatic ecology. *Aquatic Sciences* 57:255-289
- Tuomisto H (2010) A diversity of beta diversities: straightening up a concept gone awry. Part 2. Quantifying beta diversity and related phenomena. *Ecography* 33:23-45. doi:10.1111/j.1600-0587.2009.06148.x
- Tzedakis PC (2007a) Pollen records, last interglacial of Europe. In: Elias SA (ed) *Encyclopedia of Quaternary Science*, vol 3. Elsevier, Amsterdam, pp 2597-2605
- Tzedakis PC (2007b) Seven ambiguities in the Mediterranean palaeoenvironmental narrative. *Quaternary Science Reviews* 26 (17-18):2042-2066. doi:10.1016/j.quascirev.2007.03.014
- Tzedakis PC, Andrieu V, de Beaulieu JL, Birks HJB, Crowhurst S, Follieri M, Hooghiemstra H, Magri D, Reille M, Sadori L, Shackleton NJ, Wijmstra TA (2001) Establishing a terrestrial chronological framework as a basis for biostratigraphical comparisons. *Quaternary Science Reviews* 20 (16-17):1583-1592
- Tzedakis PC, Channell JET, Hodell DA, Kleiven HF, Skinner LC (2012) Determining the natural length of the current interglacial. *Nature Geoscience* 5 (2):138-141. doi:10.1038/ngeo1358
- Tzedakis PC, Drysdale RN, Margari V, Skinner LC, Menviel L, Rhodes RH, Taschetto AS, Hodell DA, Crowhurst SJ, Hellstrom JC, Fallick AE, Grimalt JO, McManus JF, Martrat B, Mokeddem Z, Parrenin F, Regattieri E, Roe K, Zanchetta G (2018) Enhanced climate instability in the North Atlantic and southern Europe during the Last Interglacial. *Nature Communications* 9 (1):4235. doi:10.1038/s41467-018-06683-3
- Tzedakis PC, Emerson BC, Hewitt GM (2013) Cryptic or mystic? Glacial tree refugia in northern Europe. *Trends in Ecology & Evolution* 28 (12):696-704. doi:10.1016/j.tree.2013.09.001
- Williams JW, Blois JL, Shuman BN (2011a) Extrinsic and intrinsic forcing of abrupt ecological change: case studies from the late Quaternary. *Journal of Ecology* 99:664-677. doi:10.1111/j.1365-2745.2011.01810.x
- Zagwijn WH (1996) An analysis of Eemian climate in western and central Europe. *Quaternary Science Reviews* 15:451-469

Figure captions




Fig. 1 Hypothetical responses of an ecosystem in the last interglacial (Eemian) in terms of biomass and fertility (modified from Birks & Birks, 2004). **a** The three phases of Eemian vegetation history, namely protocratic, mesocratic, and oligocratic plus telocratic, in response to changing temperature (outer circle). **b** Hypothetical model of compositional change (turnover) within an Eemian pollen sequence with expected patterns of turnover in each geographical region and an indication of total palynological turnover and total palynological variation expected in north (above 60°N), central (45–60°N), and south (below 45°N) Europe. The turnover axis can be, for example, an ordination axis.




Fig. 2 Map of the 47 analysed Eemian pollen sequences and individual sample scores (standard deviation units) of selected detrended canonical correspondence analysis (DCCA) axis 1 plots for 18 sequences (the remaining 29 sequences are in ESM 8). Colours of locations indicate assigned region (blue = north; green = central; salmon = south). The number at each location corresponds with sequence numbers in ESM 2. The sample scores are plotted with the oldest (bottom) at the far left and the youngest (top) at the far right. The individual sample scores are weighted averages of the response (pollen taxa) variable scores (also applies to ESM 8).




Fig. 3 Total compositional turnover in each sequence plotted against latitude estimated by **a** detrended canonical correspondence analysis (DCCA; in standard deviation units) and **b** principal curves (PC; proportional distance); **c** total inertia or variation for each sequence estimated by DCCA; and **d** the proportions of optimal partitions to total number of samples in a sequence for each region. The fitted lines in **a**, **b**, and **c** are a fitted generalised linear model with a Gaussian distribution with a second-order polynomial (grey shading is the 95% confidence interval). The DCCA-based turnover for each sequence is estimated as the range in sample scores (weighted averages of the taxon scores) within the sequence.